

First detailed analysis of multiple system V2083 Cyg

P.Zasche^{1*} and P. Svoboda² and M. Šlechta³

¹*Astronomical Institute, Faculty of Mathematics and Physics, Charles University Prague, CZ-180 00 Praha 8, V Holešovičkách 2, Czech Republic*

²*Private observatory, Výpustky 5, Brno, CZ-614 00 Czech Republic*

³*Astronomical Institute, Academy of sciences, Fričova 298, CZ-251 65, Ondřejov, Czech Republic*

Accepted ... Received ...; in original form ...

ABSTRACT

Main aim of this paper is the first detailed analysis of multiple system V2083 Cyg and to reveal its basic physical properties. The system was studied by method of the light and radial velocity curves analysis, together with the interferometric data of the visual pair obtained during a last century. There was found that the close subsystem contains two very similar stars of spectral type A7-8. Moreover, the third body is orbiting around this pair with period of about 177 years. Due to the discrepancy of total mass as derived from two methods, there arises that the third body is maybe also a binary, or some object with lower luminosity but higher mass than normal main-sequence star. Another explanation is that the Hipparcos value of parallax is incorrect and the system is much closer to the Sun.

Key words: binaries: eclipsing – binaries: visual – stars: fundamental parameters – stars: individual: V2083 Cyg.

1 INTRODUCTION

The eclipsing binaries as members of more complex multiple systems can provide us important information about their physical properties, as derived from different methods. This is the case of V2083 Cyg, which is the system, where the close components form an eclipsing binary, and the third distant body orbiting the close pair is detected as a visual component. Thanks to the combined analysis we are able to derive the radii, masses and evolutionary status of the close components and also some properties of the distant one. Such systems are still very rare and mostly lie relatively close to the Solar system. Nowadays, there are known only 33 such systems, where a close eclipsing binary is a member of a wide visual binary and we know both orbits, their mutual inclinations, ratio of periods, etc. Such a unique systems are the most suitable ones for studies of dynamical effects, the short and long-term evolution of the orbits, etc. (see e.g. Söderhjelm 1975).

2 THE SYSTEM V2083 CYG

The system V2083 Cyg (= HD 184242 = HIP 96011, RA $19^h 31^m 16.36^s$ DE $+47^\circ 28' 52.24''$, $V_{max} = 6.86$ mag) is an Algol-type eclipsing binary with its orbital period about 1.87 day. It is also a primary component of a visual double

star designated as WDS J19313+4729 in the Washington Double Star Catalog (WDS¹, Mason et al. 2001). The secondary component of this double star is about 220 mas distant and is a bit fainter. On the other hand, the magnitude difference is not very certain, because different authors list different values. The WDS catalogue itself gives 7.50 and 7.93 mag for both components.

The system is rather neglected one and there are only a few papers published. It was discovered as an eclipsing binary from the Hipparcos data (Perryman et al. 1997), which also reveal that the light curve (hereafter LC) shows two similar minima and a classical feature of an Algol-type star.

Spectral type of the system is not known very precisely nowadays. Abt (1985) presented spectral classification of the whole AB system as Am (K/H/M=A3/A8/A9), Renson et al. (1991) gives a composite spectral type as A3-A9, while the spectral type A3 was presented by Cannon & Pickering (1918), Ochsenbein (1980), and many others. This could indicate that the combined spectrum is composed from components of slightly different spectral types. The photometry of V2083 Cyg obtained from the Hipparcos mission gives a color index $B - V = 0.279$ mag (indicating sp.type A9, Popper 1980), while the infrared $J - H$ and $H - K$ indices, which are less influenced by interstellar reddening, as derived from the 2MASS survey give spectral types about A4 and A7, Cox (2000).

* E-mail: zasche@sirrah.troja.mff.cuni.cz

¹ <http://ad.usno.navy.mil/wds/>




Figure 1. Light curves in *B* (blue), *V* (green), and *R* (red) filters for V2083 Cyg, the solid line represents the final fit (see the text).

The visual orbit of the two components was derived by Seymour et al. (2002). They presented the orbital period of the double about 372 yr, the angular semimajor axis about 498 mas, and the eccentricity 0.16. However, as they already mentioned, the orbit is still only a preliminary one.

3 PHOTOMETRY AND SPECTROSCOPY

We started collecting the photometric data of the system in April 2008. In total there are 31 nights of observations, but for the light-curve analysis we used only 27 nights of observations obtained from April 2008 to September 2009 and carried out with the same telescope and detector at the private observatory by one of the authors (PS). Owing to high brightness of the target, there was used only a small 34-mm refractor at the Private observatory in Brno, Czech Republic, using the SBIG ST-7XME CCD camera and standard *BVR* filters by the specification by Bessell (1990). All the measurements were processed by the software C-MUNIPACK², which is based on aperture photometry and using the standard DaoPhot routines (Tody 1993). The other nights were used only for deriving the precise times of minima for the system.

Besides our new observations, there were also used the photometric data obtained within the SuperWASP survey (Pollacco et al. 2006). However, these data are not of enough quality to be used for the LC analysis. Hence, we made use of the SWASP photometry only for deriving the minima times of V2083 Cyg for a prospective period analysis. For all of the minima the Kwee - van Woerden method was used (Kwee & van Woerden 1956), and all of them are given in Table A1. The linear ephemeris are as follows: $HJD = 2448501.1237 + 1.867493429 \cdot E$.

The CCD spectra were obtained at Ondřejov observatory, Czech Republic, using the 2.0-m telescope equipped with a SITE-005 800×2000 CCD detector. These spectra cover a wavelength region 626 – 676 nm. All of them were secured between April 2010 and May 2011 and have a linear dispersion of about 17 Å/mm. Their S/N values range typically between 100 and 200.

For all of the spectra used the wavelength calibration was made via a ThAr comparison spectra obtained before and after the stellar spectra itself. The flatfields were taken in the beginning and end of the night and their means were used for the reduction. After then, the radial




Figure 2. Radial velocity curves of V2083 Cyg for the primary (black points), and the secondary (red points), the lines represent the final fit (see the text and Table 2).

velocities (hereafter RV) were obtained with the program SPEFO (Horn et al. 1996), with using the zero point correction via measuring the telluric lines. In total 19 spectra were obtained this way. Moreover, also two Elodie spectra (Moultaka et al. 2004) obtained in 1999 were added for the analysis.

A list of derived radial velocities from all of the available spectral observations is written in Table 1. In the last column the reference Elodie or Ondřejov is given. For all of the spectra we also tried to identify the third component lines, however these radial velocities are rather uncertain and affected by relatively large errors (see below Sections 5 and 6).

4 LC AND RV ANALYSIS

The complete LC (in *BVR* filters) and RV curves were analyzed simultaneously, using the program PHOEBE (Prša & Zwitter 2005), which is based on the Wilson-Devinney algorithm (Wilson & Devinney 1971). The derived quantities are as follows: the semi-major axis a , the mass ratio $q = M_2/M_1$, the systemic velocity γ , the secondary temperature T_2 , the inclination i , the luminosities L_i , the gravity darkening coefficients g_i , the limb darkening coefficients x_i , the albedo coefficients A_i , and the synchronicity parameters F_i . The limb darkening was approximated via linear cosine law, and the values of x_i were interpolated from van Hamme's tables, see van Hamme (1993).

For the whole analysis, we followed this procedure: at the beginning, we fixed the temperature of the primary component at $T_1 = 7930$ K (corresponding to spectral type A7, Cox 2000). We were trying to find the best LC+RV fit according to the lowest value of rms. There was a solution reached, but this one was unacceptable due to the fact that resulting values of M_1, M_2, L_1, L_2, T_1 , and T_2 are in contradiction with each other. In particular, the resulting spectral types as derived from M, L and T differ significantly between each other. For this reason, we tried a different starting value of T_1 . With this method we were changing the temperature T_1 in the range from 8520 K to 7020 K (spectral types A3 to F0) and trying to find a consistent solution. For all of these attempts, the value of T_1 remained fixed.

Our final parameters as derived from the LC+RV fit are given in Table 2. The plot of the LC is shown in Fig. 1, while the RV curves with the fits are given in Fig. 2. The value of eccentricity was fixed at 0. For the discussion about

² see <http://c-munipack.sourceforge.net/>




Figure 3. Visual orbit of V2083 Cyg as displayed on the sky. The individual observations (black points) are connected with their theoretical positions (red points). The dashed line represents the line of the nodes, while the dotted one the line of the apsides. Parameters of the fit are given in Table 3.

the physical parameters of the components (eclipsing ones and also the third one), see Section 6.

For the whole computation process, the values of albedos A_i and the values of gravity darkening coefficients were set at their appropriate values ($A_i = 1$ or 0.5 , and $g_i = 1$ or 0.32) according to the component's temperatures ($T_i < 7200$ K, or $T_i > 7200$ K). Another problematic issue were the values of F_i , which tended to decrease down to 0 for both components for each of the T_1 values. These dropped down very quickly after a few steps of iterations. Due to this reason we tried a different approach. From the spectra of the system we estimated the values $v \sin i$, which were used to derive the values of F_i for both components. Therefore, the values of F_i as given in Table 2 are not derived from the combined LC&RV analysis but from the spectra.

The fitting process with PHOEBE was carried out assuming three luminosities. Besides the luminosities of primary and secondary component of the eclipsing binary pair also the additional third light L_3 was considered. This luminosity corresponds to the visual component B and is presented in the combined light for all the time (the two visual components are too close). From this value one can speculate about some physical parameters of the third body in the system, see below Section 6.

5 VISUAL ORBIT

The close eclipsing pair is orbiting around a common barycenter with the third distant component of the system. Recent precise interferometric observations are to be used for determining the parameters of this visual orbit. Since its discovery as a double star by Aitken (1904) there were obtained 61 astrometric observations of the double (i.e. position angle and separation). We took these data from the

Table 1. Radial velocities of V2083 Cyg as derived from the spectra from the ELODIE archive, and from the Ondřejov observatory.

HJD - 2400000	RV1	RV2	RV3	Ref.
51405.3658	95.594	-81.861		Elodie
51407.4156	129.872	-117.06		Elodie
55316.5290	20.920	2.771	-16.272	OND
55380.5210	-122.394	142.771	-16.336	OND
55385.3570	122.946	-99.062	-16.775	OND
55385.5530	56.001	-29.860	-18.597	OND
55386.3640	-73.970	99.094	-14.986	OND
55386.5650	-.442	25.184	-17.640	OND
55405.3670	55.349	-26.402	-16.327	OND
55425.3970	-117.384	141.598	-16.145	OND
55496.2990	-121.010	140.992	-16.273	OND
55496.4630	-99.677	124.485	-16.277	OND
55497.2890	141.451	-119.863	-15.212	OND
55622.5420	116.769	-96.782	-15.048	OND
55622.6370	89.474	-66.160	-14.284	OND
55671.4500	-18.008	36.024	-14.574	OND
55671.5840	-72.089	92.471	-14.950	OND
55671.6310	-86.135	105.183	-14.784	OND
55689.5750	144.545	-122.291	-16.380	OND
55692.3700	-119.493	143.646	-15.231	OND
55692.5540	-98.435	122.422	-15.080	OND

Table 2. The final LC and RV parameters of V2083 Cyg.

Parameter	Value	Parameter	Value
a [$R_\odot$]	9.57 ± 0.15	L_1 (B) [%]	26.7 ± 0.7
$q = M_2/M_1$	0.97 ± 0.07	L_2 (B) [%]	34.9 ± 0.9
γ [km/s]	10.78 ± 0.68	L_3 (B) [%]	38.4 ± 0.8
T_1 [K]	7630 (fixed)	L_1 (V) [%]	26.7 ± 0.6
T_2 [K]	7623 ± 45	L_2 (V) [%]	34.7 ± 0.9
e	0 (fixed)	L_3 (V) [%]	38.6 ± 0.8
i [deg]	80.47 ± 1.60	L_1 (R) [%]	26.4 ± 0.6
$x_1 = x_2$ (B)	0.412	L_2 (R) [%]	34.2 ± 0.9
$x_1 = x_2$ (V)	0.356	L_3 (R) [%]	39.4 ± 1.0
$x_1 = x_2$ (R)	0.356	Derived physical quantities:	
$g_1 = g_2$	1.000 (fixed)	R_1 [$R_\odot$]	2.12 ± 0.17
$A_1 = A_2$	1.000 (fixed)	R_2 [$R_\odot$]	2.45 ± 0.20
F_1	0.81 ± 0.13	M_1 [$M_\odot$]	1.71 ± 0.11
F_2	0.84 ± 0.11	M_2 [$M_\odot$]	1.66 ± 0.09

WDS database. The very last observation was obtained in 2009.

Since its discovery, the position angle of the pair changed of about 88° . Thanks to this movement, the orbit of the pair around a barycenter was derived. The orbit was published by Seymour et al. (2002), who computed its orbital period of about 372 yr. However, since this most recent study there were published three new interferometric observations, so we decided to perform a new analysis with the complete data set.

Our new computation led to the visual orbit parameters given in Table 3. For the computation we used a following approach. Starting with the orbital parameters as published by Seymour et al. (2002), the final fit reached very different solution. Moreover, there were found several different minima in the parameter space as derived from this astrometric data set. Some minimum was found with very long orbital




Figure 4. Plot of period-mass and period-rms relation. The dashed lines represent our final fit. For details see the text.

periods, but this solution seems to be less probable due to the poor coverage of the data. The most significant minimum (the deepest one) was found near the period of 177.4 years. However, we would like to emphasize that the orbital solution is still a preliminary one. New precise observations secured every year would be very welcome for derivation of the orbital parameters with higher conclusiveness and especially for setting the more solid constraints on p_3 and a values. These are the most important for a discussion about the nature of the third component (see Section 6 below).

In Fig. 4 there is a plot of total mass versus period, as well as the rms of the particular fit versus period. For our final solution reached (minimum rms with $p_3 = 64778.357$ day) the value of total mass was computed (using the Hipparcos parallax) - these are shown as dashed lines in Fig. 4. The relation between the two vertical axis (parallax and total mass) is defined via a third Kepler's law using our final solution. As one can see from the bold line of mass-period relation, the total mass as derived from our final solution is close to the minimal mass in this period range (the uncertainty of Hipparcos parallax $\pi_{HIP} = 4.32 \pm 0.57$ is shown as a grey area). Of course, this analysis is very sensitive to the input weighting scheme. The individual weights of the data points were set equal to each other, because for most of the observations the sigma or some other error estimations are missing. No minimum of rms near a period of 372 years as proposed by Seymour et al. (2002) is presented. One can ask, why such a different solution was reached when using only three new interferometric observations. The main reason (maybe besides different weighting) is that these three new measurements provide a strong constraints on the fit. This is due to the fact that the position angle between our most recent data and those from Seymour et al. (2002) changed about 20° , which is about $1/4$ of the total position angle range covered. All of these calculations (e.g. the Kepler's law) were using the set of recommended values of fundamental parameters as proposed by Harmanec & Prša (2011).

On the other hand, we also tried to compute the predicted change of the third-body velocities over the time span of more than 11 years covered with our spectroscopic data. Taken into account some assumptions (masses), the change

Table 3. Final parameters of the long orbit.

Parameter	Seymour et al. (2002)	This work
p_3 [day]	135869	64778 ± 427
p_3 [yr]	372	177.4 ± 1.2
a [mas]	498	291.9 ± 1.4
T_0	2438395	2400006 ± 375
Ω [deg]	73.6	174.54 ± 2.9
ω [deg]	189	334.89 ± 5.3
i [deg]	64	48.73 ± 3.6
e	0.16	0.471 ± 0.018

in velocity resulted in more than 20 km/s. Such a large velocity difference should be easily detectable in our RV_3 data. Unfortunately, we were not able to identify the third-component lines in the ELODIE spectra and in newer data from Ondřejov there is no such difference, hence we could only speculate about our findings. The reason could be either different masses or much longer orbital period. Another explanation is an incorrect identification of the third-body lines in the spectra.

6 PHYSICAL PARAMETERS

Taken into account all results as presented above, one can make a picture of the system, its geometry and orientation in space. From the combined LC and RV analysis there resulted that both eclipsing components are probably main sequence stars, located well within their respective Roche lobes. According to their masses and temperatures, it seems like their individual spectral types are probably of A7 and A8 (e.g. Popper 1980, Harmanec 1988, or Andersen 1991) for primary and secondary, respectively. However, according to their luminosities, it seems like the stars are of slightly earlier spectral type (about A5).

Another task was to derive the value of the third light L_3 from the LC solution and to obtain a magnitude difference between the two visual components. This value resulted in about 0.49 mag, which is in rough agreement with the $\Delta m = 0.43$ mag value as presented in the WDS catalogue.

A discussion about the third body is still difficult due to some aspects of the problem. The most problematic issue is still the uncertainty of the Hipparcos value of parallax. The relatively high error of about 13 % could lead to distances in wide range from 204 to 267 pc. Thanks to this uncertainty also the value of total mass as computed from the visual orbit (see Table 3) lie in between 6.54 and 15.41 $M_{\odot}$ with the mean value of 9.81 $M_{\odot}$.

Subtracting masses of both eclipsing components, we obtain an interesting result of mass of the third body about 6.44 $M_{\odot}$ (with upper and lower limits about 12.24 and 2.97). Such a massive third body easily cannot be a main sequence A star as predicted from the Δm value. One possible explanation of this discrepancy is that this component is also a double star. If we speculate about two identical stars, then such stars have to be of only slightly later spectral type than the eclipsing components. (because of the total mass). Assuming two F0 stars, we can hardly satisfy the magnitude difference between the components. However, this explana-

tion is still questionable because the third lines in the spectra do not show a doubleness profile.

To solve this discrepancy we also tried to use the program KOREL (Hadrava 2004) for disentangling of the spectra taken at Ondřejov observatory. However, it was also not able to solve the problem. The final parameters on one hand confirmed our findings about the LC+RV solution (mass ratio q from KOREL resulted in about 0.993), but on the other hand it also results with the value of mass ratio $q_3 = M_3/M_{12} > 1$. This would indicate that the third body is more massive than the eclipsing pair, but also the less luminous one. Solving the problem of its lower luminosity and higher mass with introducing a degenerate object is a highly speculative solution. Hence, the nature of the third body still remains an open question. The KOREL radial velocities of the third body were also used and these are the values presented in Table 1 in the *RV3* column.

7 DISCUSSION AND CONCLUSIONS

The multiple system V2083 Cyg is still rather neglected one and this is the first detailed analysis of it. The components of an eclipsing binary are of spectral type A, are well-detached, with no evidence of circumstellar matter, emission in the spectra, etc. This close pair is also orbiting around a common barycenter with the third component with period about 177 years. The mutual inclination of the two orbits is 31.8° , therefore we can only speculate about a common origin of the system.

A nature of the third component is still rather problematic to derive. From the combined LC and RV analysis there results that the third body is slightly less luminous than the eclipsing pair. But the Hipparcos parallax indicates a higher total mass of the system than computed from all components' masses. A possible explanation is that the value of Hipparcos parallax is underestimated and a real distance of V2083 Cyg is lower (even outside of the error bars of the Hipparcos data). This would not be an exceptional case, because for some systems the Hipparcos data yielded incorrect parallax due to the presence of a close visual companion (e.g. Docobo et al. 2008). Another possible explanation is that the body is also a binary, but there are also some problems with this explanation (luminosity and the spectral lines of such body). For that reason, new more detailed observations are still very welcome.

However, if the hypothesis of binarity of the third component is proved, it will shift the triple to quadruple. On one hand, such systems of higher multiplicity are of big interest, on the other hand we also deal with very incomplete statistics of them among the stars (see e.g. Eggleton & Tokovinin 2008, and Eggleton 2009).

ACKNOWLEDGMENTS

We thank the "SWASP" team for making all of the observations easily public available. Mr. Robert Uhlař is acknowledged for sending us his photometric data and Prof. Petr Harmanec for a useful discussion and valuable advices. We also thank an anonymous referee for his/her helpful and

critical suggestions. We also acknowledge the use of spectrograms from the public archives of the Elodie spectrograph of the Haute Provence Observatory. We are obliged to our colleagues who took spectra of V2083 Cyg in Ondřejov observatory (M. Wolf, P. Chadima, and J.A. Nemravová). This work was supported by the Czech Science Foundation grant no. P209/10/0715 and also by the Research Programme MSM0021620860 of the Czech Ministry of Education. This research has made use of the Washington Double Star Catalog maintained at the U.S. Naval Observatory, the SIMBAD database, operated at CDS, Strasbourg, France, and of NASA's Astrophysics Data System Bibliographic Services.

REFERENCES

- Abt, H. A. 1985, *ApJS*, 59, 95
- Aitken, R. G. 1904, *LicOB*, 3, 6
- Andersen, J. 1991, *A&ARv*, 3, 91
- Bessell, M. S. 1990, *PASP*, 102, 1181
- Cannon, A. J., Pickering, E. C. 1918, *AnHar*, 91, 1
- Cox, A. N. 2000, *Allen's Astrophysical Quantities*
- Docobo, J. A., Tamazian, V. S., Andrade, M., Melikian, N. D., & Karapetian, A. A. 2008, *AJ* 136, 890
- Eggleton, P. P., Tokovinin, A. A. 2008, *MNRAS*, 389, 869
- Eggleton, P. P. 2009, *MNRAS*, 399, 1471
- Hadrava, P. 2004, *PAICz*, 92, 15
- van Hamme, W. 1993, *AJ*, 106, 2096
- Harmanec, P. 1988, *BAICz*, 39, 329
- Harmanec, P., & Prša, A. 2011, *PASP*, 123, 976
- Horn, J., Kubát, J., Harmanec, P., Koubský, P., Hadrava, P., Šimon, V., Štefl, S., Škoda, P. 1996, *A&A*, 309, 521
- Kwee, K.K., van Woerden, H. 1956, *BAN*, 12, 327
- Mason, B. D., Wycoff, G. L., Hartkopf, W. I., Douglass, G. G., Worley, C. E. 2001, *AJ*, 122, 3466
- Moultaka, J., Ilovaisky, S. A., Prugniel, P., Soubiran, C. 2004, *PASP*, 116, 693
- Ochsenbein, F. 1980, *BICDS*, 19, 74
- Perryman, M. A. C., Lindegren, L., Kovalevsky, J., et al. 1997, *A&A*, 323, L49
- Pollacco, D. L., et al. 2006, *PASP*, 118, 1407
- Popper, D. M. 1980, *ARA&A*, 18, 115
- Prša, A., Zwitter, T. 2005, *ApJ*, 628, 426
- Renson, P., Gerbaldi, M., Catalano, F. A. 1991, *A&AS*, 89, 429
- Seymour, D. M., Mason, B. D., Hartkopf, W. I., Wycoff, G. L. 2002, *AJ*, 123, 1023
- Söderhjelm, S. 1975, *A&A*, 42, 229
- Tody, D. 1993, "IRAF in the Nineties" in *Astronomical Data Analysis Software and Systems II*, A.S.P. Conference Ser., Vol 52, eds. R.J. Hanisch, R.J.V. Brissenden, J. Barnes, 173.
- Wilson, R. E., Devinney, E. J. 1971, *ApJ*, 166, 605

APPENDIX A:

Table A1. Heliocentric minima of V2083 Cyg.

HJD - 2400000	Error	Type	Filter	Observer
54609.69354	0.00179	prim	-	SWASP
54638.63794	0.00035	sec	-	SWASP
54639.57227	0.00042	prim	-	SWASP
54652.64554	0.00089	prim	-	SWASP
54668.51720	0.00046	sec	-	SWASP
54669.44747	0.00046	prim	-	SWASP
54683.46156	0.00027	sec	-	SWASP
54684.39467	0.00095	prim	-	SWASP
54994.39887	0.00091	prim	B	PS
54994.39888	0.00113	prim	V	PS
54994.39948	0.00128	prim	R	PS
55049.49292	0.00110	sec	B	PS
55049.49015	0.00104	sec	R	PS
55049.49055	0.00062	sec	V	PS
55051.35585	0.00081	sec	B	PS
55051.35567	0.00078	sec	V	PS
55051.35645	0.00107	sec	R	PS
55064.43157	0.00091	sec	B	PS
55064.43162	0.00057	sec	V	PS
55064.43102	0.00064	sec	R	PS
55076.57570	0.00116	prim	B	PS
55076.57197	0.00108	prim	V	PS
55076.57194	0.00165	prim	R	PS
55093.37749	0.00087	prim	B	PS
55093.37848	0.00068	prim	V	PS
55093.37624	0.00063	prim	R	PS
55374.43431	0.00021	sec	I	RU
55429.52450	0.00028	prim	I	RU
55740.46289	0.00056	sec	R	PS
55797.42284	0.00043	prim	I	RU

Note: PS - Petr Svoboda, RU - Robert Uhlář